

OMEGA-CATEGORIES AND CHAIN COMPLEXES

RICHARD STEINER

(communicated by Ronald Brown)

Abstract

There are several ways to construct omega-categories from combinatorial objects such as pasting schemes or parity complexes. We make these constructions into a functor on a category of chain complexes with additional structure, which we call augmented directed complexes. This functor from augmented directed complexes to omega-categories has a left adjoint, and the adjunction restricts to an equivalence on a category of augmented directed complexes with good bases. The omega-categories equivalent to augmented directed complexes with good bases include the omega-categories associated to globes, simplexes and cubes; thus the morphisms between these omega-categories are determined by morphisms between chain complexes. It follows that the entire theory of omega-categories can be expressed in terms of chain complexes; in particular we describe the biclosed monoidal structure on omega-categories and calculate some internal homomorphism objects.

1. Introduction

This paper is a contribution to the theory of strict ω -categories. In the past, ω -categories have been constructed from combinatorial structures such as pasting schemes, parity complexes or directed complexes; see Johnson [6], Power [9], Steiner [10] and Street [12]. The constructions are not really functorial, although there are ways to produce non-obvious morphisms; see Crans-Steiner [4]. The combinatorial structures can be regarded as bases for chain complexes, as observed by Kapranov and Voevodsky in [7]. We will reinterpret the constructions as a functor on a category of chain complexes with additional structure called augmented directed complexes, and we will show that the functor has a left adjoint. The adjunction is related to the well-known equivalence between the categories of chain complexes and of ω -categories in the category of abelian groups; see Brown-Higgins [3] for instance.

In the earlier treatments, it is shown that combinatorial structures which are in a suitable sense loop-free produce free ω -categories. The functorial version of this result says that the adjunction restricts to equivalences between certain pairs of full

Received March 11, 2004, revised April 27, 2004; published on May 12, 2004.

2000 Mathematics Subject Classification: 18D05.

Key words and phrases: Omega-category, augmented directed complex.

© 2004, Richard Steiner. Permission to copy for private use granted.

subcategories. The augmented directed complexes concerned are free chain complexes with good bases, and the corresponding ω -categories have good sets of generators. These ω -categories include those associated to globes, simplexes and cubes, which determine the entire theory of ω -categories (see Al-Agl-Brown-Steiner [1], Al-Agl-Steiner [2] and Street [11]); thus the theory of ω -categories can be described in terms of chain complexes. In particular the biclosed monoidal structure on ω -categories can be described in terms of chain complexes, and we will calculate some internal homomorphism objects. Homomorphisms between these ω -categories have been studied by Gaucher in his work on higher-dimensional automata [5].

The adjunction between ω -categories and augmented directed complexes is described in Section 2, bases for augmented directed complexes are described in Section 3, generating sets for ω -categories are described in Section 4, the equivalences between subcategories are described in Section 5, relations with earlier work are described in Section 6, and the applications to the theory of ω -categories are described in Section 7.

2. The adjunction

In this section we describe the adjunction between ω -categories and augmented directed complexes. First we define the categories involved. For ω -categories we use the following description and notation.

Definition 2.1. An ω -category is a set with unary source and target operators d_0^- , d_0^+ , d_1^- , d_1^+ , \dots and not everywhere defined binary composition operators

$$(x, y) \mapsto x \#_0 y, (x, y) \mapsto x \#_1 y, \dots$$

such that the following hold:

- (i) $x \#_n y$ is defined if and only if $d_n^+ x = d_n^- y$;
- (ii) for every x there exists n such that $d_n^- x = d_n^+ x = x$;
- (iii) for any x ,

$$d_m^\beta d_n^\alpha x = \begin{cases} d_m^\beta x & \text{if } m < n, \\ d_n^\alpha x & \text{if } m \geq n; \end{cases}$$

- (iv) for any x ,

$$d_n^- x \#_n x = x \#_n d_n^+ x = x;$$

- (v) if $x \#_n y$ is defined then

$$d_m^\alpha (x \#_n y) = d_m^\alpha x = d_m^\alpha y = d_m^\alpha x \#_n d_m^\alpha y \text{ for } m < n,$$

$$d_n^- (x \#_n y) = d_n^- x,$$

$$d_n^+ (x \#_n y) = d_n^+ y,$$

$$d_m^\alpha (x \#_n y) = d_m^\alpha x \#_n d_m^\alpha y \text{ for } m > n;$$

- (vi) for any n ,

$$(x \#_n y) \#_n z = x \#_n (y \#_n z)$$

if either side is defined;

(vii) if $m < n$ then

$$(x \#_n y) \#_m (x' \#_n y') = (x \#_m x') \#_n (y \#_m y')$$

when the left side is defined.

A morphism of ω -categories is a function commuting with the source, target and composition operators. The category of ω -categories is denoted $\omega\text{-cat}$.

Let C be an ω -category. It generates an ω -category object C' in the category of abelian groups, which is equivalent to a nonnegatively graded chain complex K by [3]. On C' there are two pieces of additional structure: the homomorphism $C' \rightarrow \mathbf{Z}$ induced by the morphism from C to the one-element ω -category, and the submonoid generated by the image of C (a submonoid of an abelian group is a subset containing zero and closed under addition). Correspondingly, it turns out that the chain complex K is augmented and has a distinguished submonoid in each chain group. A chain complex with this kind of additional structure will be called an augmented directed complex.

Definition 2.2. An *augmented directed complex* K is an augmented chain complex $(K_n, \partial, \varepsilon)$ of abelian groups concentrated in nonnegative dimensions, together with a distinguished submonoid K_n^* of the chain group K_n for each n . A morphism of augmented directed complexes from K to L is an augmentation-preserving chain map $f: K \rightarrow L$ such that $f(K_n^*) \subset L_n^*$ for each n . The category of augmented directed complexes is denoted **ADC**.

Similar structures have been studied by Patchkoria in [8].

In order to construct the functor from $\omega\text{-cat}$ to **ADC**, we first recall the standard filtration for an ω -category.

Proposition 2.3. *Let C be an ω -category. Then C is the union of an increasing sequence of sub- ω -categories denoted $C_0 \subset C_1 \subset \dots$, where*

$$C_n = d_n^- C = d_n^+ C = \{x \in C : d_n^- x = x\} = \{x \in C : d_n^+ x = x\}.$$

Proof. The four definitions for C_n are consistent because $d_n^\beta d_n^\alpha = d_n^\alpha$. The C_n are sub- ω -categories by Definition 2.1(iii) and (vi). They form an increasing sequence because $d_n^\alpha = d_{n+1}^\beta d_n^\alpha$, and C is the union of the sub- ω -categories by Definition 2.1(ii). \square

The filtration of Proposition 2.3 is such that the composition operators $\#_m$ are trivial in C_n for $m \geq n$; thus C_n is an n -category.

We now use the filtration to construct a functor from $\omega\text{-cat}$ to **ADC**.

Definition 2.4. The functor $\lambda: \omega\text{-cat} \rightarrow \mathbf{ADC}$ is defined as follows. Let C be an ω -category. Then the chain group $(\lambda C)_n$ for $n \geq 0$ is generated by elements $[x]_n$ for $x \in C_n$ subject to relations

$$[x \#_m y]_n = [x]_n + [y]_n \text{ for } m < n,$$

the boundary homomorphism $\partial: (\lambda C)_{n+1} \rightarrow (\lambda C)_n$ for $n \geq 0$ is given by

$$\partial[x]_{n+1} = [d_n^+ x]_n - [d_n^- x]_n,$$

the augmentation $\varepsilon: (\lambda C)_0 \rightarrow \mathbf{Z}$ is given by

$$\varepsilon[x]_0 = 1,$$

the distinguished submonoid $(\lambda C)_n^*$ is the submonoid generated by the elements $[x]_n$.

To justify this definition, we make the following observations. First, if $x \in C_{n+1}$ then the $d_n^\alpha x$ are members of C_n , so the difference $[d_n^+ x]_n - [d_n^- x]_n$ is a member of $(\lambda C)_n$. If $x \#_n y$ is a composite in C_{n+1} then

$$[d_n^+(x \#_n y)]_n - [d_n^-(x \#_n y)]_n = [d_n^+ y]_n - [d_n^- x]_n = ([d_n^+ x]_n - [d_n^- x]_n) + ([d_n^+ y]_n - [d_n^- y]_n)$$

because $d_n^+ x = d_n^- y$, and if $x \#_m y$ is a composite in C_{n+1} with $m < n$ then

$$\begin{aligned} [d_n^+(x \#_m y)]_n - [d_n^-(x \#_m y)]_n &= [d_n^+ x \#_m d_n^+ y]_n - [d_n^- x \#_m d_n^- y]_n \\ &= ([d_n^+ x]_n + [d_n^+ y]_n) - ([d_n^- x]_n + [d_n^- y]_n) \\ &= ([d_n^+ x]_n - [d_n^- x]_n) + ([d_n^+ y]_n - [d_n^- y]_n); \end{aligned}$$

therefore $\partial: (\lambda C)_{n+1} \rightarrow (\lambda C)_n$ is a well-defined homomorphism. There are no relations on $(\lambda C)_0$, so $\varepsilon: (\lambda C)_0 \rightarrow \mathbf{Z}$ is a well-defined homomorphism. The composites $\partial\partial: (\lambda C)_{n+2} \rightarrow (\lambda C)_n$ are trivial because $d_n^\alpha d_{n+1}^- = d_n^\alpha d_{n+1}^+$, and the composite $\varepsilon\partial: (\lambda C)_1 \rightarrow \mathbf{Z}$ is obviously trivial. Finally, λ is obviously functorial.

We make the following observation.

Proposition 2.5. *If C is an ω -category and $x \in C_m$ with $m < n$, then $[x]_n = 0$ in $(\lambda C)_n$.*

Proof. This holds because

$$[x]_n = [x \#_m d_m^+ x]_n = [x \#_m x]_n = [x]_n + [x]_n.$$

□

We will now go from chain complexes to ω -categories. We first define a functor μ from arbitrary chain complexes to ω -categories in the category of abelian groups. It is the equivalence given implicitly in [3], and it may also be regarded an additive version of Street's construction [12].

Definition 2.6. The functor μ from chain complexes to ω -categories in the category of abelian groups is defined as follows. Let K be a chain complex. Then μK is the abelian group of double sequences

$$x = (x_0^-, x_0^+, x_1^-, x_1^+, \dots)$$

such that

$$\begin{aligned} x_n^- &\in K_n \text{ and } x_n^+ \in K_n, \\ x_n^- &= x_n^+ = 0 \text{ for all but finitely many values of } n, \\ x_n^+ - x_n^- &= \partial x_{n+1}^- = \partial x_{n+1}^+ \text{ for } n \geq 0; \end{aligned}$$

if $x \in \mu K$ then

$$d_n^\alpha x = (x_0^-, x_0^+, \dots, x_{n-1}^-, x_{n-1}^+, x_n^\alpha, x_n^\alpha, 0, 0, \dots);$$

if $d_n^+x = d_n^-y = z$, say, in μK then

$$\begin{aligned} x \#_n y &= x - z + y \\ &= (x_0^-, y_0^+, \dots, x_n^-, y_n^+, x_{n+1}^- + y_{n+1}^-, x_{n+1}^+ + y_{n+1}^+, \dots). \end{aligned}$$

It is straightforward to check that μ is a well-defined functor. We extend the filtration of μK given in Proposition 2.3 by writing $(\mu K)_{-1} = 0$, and we get the following result.

Proposition 2.7. *Let K be a chain complex and let n be a nonnegative integer. Then $x_n^- = x_n^+$ for $x \in (\mu K)_n$, and the homomorphism $x \mapsto x_n^\alpha: (\mu K)_n \rightarrow K_n$ fits into a natural split short exact sequence*

$$0 \rightarrow (\mu K)_{n-1} \rightarrow (\mu K)_n \rightarrow K_n \rightarrow 0.$$

Proof. If $x \in (\mu K)_n$ then $x = d_n^\alpha x$, so $x_n^- = x_n^+$. For $x \in \mu K$, one finds that $x \in (\mu K)_n$ if and only if $x_m^- = x_m^+ = 0$ for all $m > n$, and it follows that the homomorphism $x \mapsto x_n^\alpha: (\mu K)_n \rightarrow K_n$ has kernel $(\mu K)_{n-1}$. This means that there is a natural exact sequence $0 \rightarrow (\mu K)_{n-1} \rightarrow (\mu K)_n \rightarrow K_n$. To complete the proof we need a natural splitting homomorphism $K_n \rightarrow (\mu K)_n$. For $n = 0$ we use the homomorphism

$$x \mapsto (x, x, 0, 0, \dots),$$

and for $n > 0$ we can use the homomorphism

$$x \mapsto (0, 0, \dots, 0, \partial x, x, x, 0, 0, \dots).$$

□

We now use the additional structure on an augmented directed complex to define a functor from **ADC** to $\omega\text{-cat}$.

Definition 2.8. The functor $\nu: \mathbf{ADC} \rightarrow \omega\text{-cat}$ is defined as follows. Let K be an augmented directed complex. Then νK is the sub- ω -category of μK consisting of the elements

$$(x_0^-, x_0^+, x_1^-, x_1^+, \dots)$$

such that

$$\begin{aligned} x_n^- \in K_n^* \text{ and } x_n^+ \in K_n^* \text{ for all } n, \\ \varepsilon x_0^- = \varepsilon x_0^+ = 1. \end{aligned}$$

It is straightforward to check that Definition 2.8 gives a well-defined functor from augmented directed complexes to ω -categories: if K is an augmented directed complex then νK is a subset of μK closed under the ω -category operations. Obviously νK is not a subgroup of μK .

Finally in this section, we show that λ is left adjoint to ν . First we describe the unit of the adjunction.

Definition 2.9. The natural transformation $\eta: C \rightarrow \nu\lambda C$ for an ω -category C is defined by the formula

$$\eta x = ([d_0^- x]_0, [d_0^+ x]_0, [d_1^- x]_1, [d_1^+ x]_1, \dots).$$

In order to justify this definition, we first show that $\eta x \in \nu\lambda C$ for $x \in C$. Indeed, it is clear that $[d_n^\alpha x]_n \in (\lambda C)_n^*$ and $\varepsilon[d_0^\alpha x]_0 = 1$, and we also have

$$\partial[d_{n+1}^\alpha x]_{n+1} = [d_n^+ d_{n+1}^\alpha x]_n - [d_n^- d_{n+1}^\alpha x]_n = [d_n^+ x]_n - [d_n^- x]_n,$$

so it suffices to show that $[d_n^\alpha x]_n = 0$ for n sufficiently large. But by Proposition 2.3 there exists p such that $d_n^\alpha x \in C_p$ for all n , and it then follows from Proposition 2.5 that $[d_n^\alpha x]_n = 0$ for $n > p$.

We must also show that $\eta: C \rightarrow \nu\lambda C$ is a morphism of ω -categories. But we get $\eta d_n^\alpha x = d_n^\alpha \eta x$ because $[d_m^\beta d_n^\alpha x]_m = [d_m^\beta x]_m$ for $m < n$, because $[d_n^\beta d_n^\alpha x]_n = [d_n^\alpha x]_n$, and because $[d_m^\beta d_n^\alpha x]_m = [d_n^\alpha x]_m = 0$ for $m > n$ by Proposition 2.5. We also get $\eta(x \#_n y) = \eta x \#_n \eta y$ because $[d_m^- (x \#_n y)]_m = [d_m^- x]_m$ and $[d_m^+ (x \#_n y)]_m = [d_m^+ y]_m$ for $m \leq n$ and because $[d_m^\alpha (x \#_n y)]_m = [d_m^\alpha x \#_n d_m^\alpha y]_m = [d_m^\alpha x]_m + [d_m^\alpha y]_m$ for $m > n$. It follows that η is a morphism of ω -categories, and it is clear that η is natural.

Next we describe the counit. We denote this by π rather than ε , in order to avoid confusion with augmentations.

Definition 2.10. The natural transformation $\pi: \lambda\nu K \rightarrow K$ is defined for an augmented directed complex K by the formula

$$\pi[x]_n = x_n^- = x_n^+.$$

To justify this definition, observe first that $x_n^- = x_n^+$ for $x \in (\nu K)_n$ by Proposition 2.7. The formula $\pi[x]_n = x_n^- = x_n^+$ gives a well-defined homomorphism $\pi: (\lambda\nu K)_n \rightarrow K_n$ because if $m < n$ then $(x \#_m y)_n^\alpha = x_n^\alpha + y_n^\alpha$. We get a chain map because

$$\partial\pi[x]_{n+1} = \partial x_{n+1}^\alpha = x_n^+ - x_n^- = (d_n^+ x)_n^+ - (d_n^- x)_n^- = \pi([d_n^+ x]_n - [d_n^- x]_n) = \pi\partial[x]_{n+1}.$$

This chain map is augmentation-preserving because $\varepsilon\pi[x]_0 = \varepsilon x_0^\alpha = 1 = \varepsilon[x]_0$ for $x \in (\nu K)_0$, and we get $\pi(\lambda\nu K)_n^* \subset K_n^*$ because $x_n^\alpha \in K_n^*$ for $x \in \nu K$. Therefore $\pi: \lambda\nu K \rightarrow K$ is a morphism of augmented directed complexes. It is clearly natural.

The main theorem is now as follows.

Theorem 2.11. *The functors $\lambda: \omega\text{-cat} \rightarrow \mathbf{ADC}$ and $\nu: \mathbf{ADC} \rightarrow \omega\text{-cat}$ form an adjoint pair with unit $\eta: \text{id} \rightarrow \nu\lambda$ and counit $\pi: \lambda\nu \rightarrow \text{id}$.*

Proof. We must show that

$$(\pi\lambda) \circ (\lambda\eta) = \text{id}: \lambda \rightarrow \lambda$$

and

$$(\nu\pi) \circ (\eta\nu) = \text{id}: \nu \rightarrow \nu.$$

Let C be an ω -category. The generators of $(\lambda C)_n$ have the form $[x]_n$ with $x \in C_n$. For these generators, $d_n^\alpha x = x$, so

$$\begin{aligned} \pi(\lambda\eta)[x]_n &= \pi[\eta x]_n \\ &= \pi[(d_0^- x]_0, [d_0^+ x]_0, [d_1^- x]_1, [d_1^+ x]_1, \dots)]_n \\ &= [d_n^\alpha x]_n \\ &= [x]_n; \end{aligned}$$

therefore $(\pi\lambda) \circ (\lambda\eta) = \text{id}$.

Now let K be an augmented directed complex and let x be a member of νK . Then

$$\begin{aligned} (\nu\pi)\eta x &= (\nu\pi)([d_0^- x]_0, [d_0^+ x]_0, [d_1^- x]_1, [d_1^+ x]_1, \dots) \\ &= (\pi[d_0^- x]_0, \pi[d_0^+ x]_0, \pi[d_1^- x]_1, \pi[d_1^+ x]_1, \dots) \\ &= ((d_0^- x)_0^-, (d_0^+ x)_0^+, (d_1^- x)_1^-, (d_1^+ x)_1^+, \dots) \\ &= (x_0^-, x_0^+, x_1^-, x_1^+, \dots) \\ &= x; \end{aligned}$$

therefore $(\nu\pi) \circ (\eta\nu) = \text{id}$.

This completes the proof. □

3. Bases for augmented directed complexes

We will now consider augmented directed complexes with bases. We essentially recover the examples constructed in earlier treatments such as [6], [9], [10] and [12].

Definition 3.1. Let K be an augmented directed complex. A *basis* for K is a set $B \subset \bigsqcup_n K_n$ such that each K_n is a free abelian group with basis $B \cap K_n$ and each K_n^* is the submonoid of K_n generated by $B \cap K_n$.

Suppose that K is an augmented directed complex with a basis. We make K_n into a partially ordered abelian group by the rule

$$x \leq y \iff y - x \in K_n^*.$$

The basis elements in K_n can be characterised as the minimal non-zero elements in K_n^* , and it follows that K has only one basis. Note also that K_n is a lattice: any two elements x and y have a least upper bound $x \vee y$ and a greatest lower bound $x \wedge y$. If x is an element of K_{n+1} , then there are unique elements $\partial^- x, \partial^+ x \in K_n$, the negative and positive parts of ∂x , such that

$$\partial x = \partial^+ x - \partial^- x, \quad \partial^- x \wedge \partial^+ x = 0;$$

indeed, if ∂x is expressed as a linear combination of distinct basis elements, then $\partial^+ x$ is the sum of the terms with positive coefficients and $-\partial^- x$ is the sum of the terms with negative coefficients.

Let b be a basis element for K . We denote the *dimension* of b by $|b|$, so that $b \in K_{|b|}$. We then define elements $\langle b \rangle_n^-$ and $\langle b \rangle_n^+$ in K_n by downward recursion as follows:

$$\langle b \rangle_n^\alpha = \begin{cases} 0 & \text{for } n > |b|, \\ b & \text{for } n = |b|, \\ \partial^\alpha \langle b \rangle_{n+1}^\alpha & \text{for } n < |b|. \end{cases}$$

It is straightforward to check that this produces an element $\langle b \rangle$ of μK ; in fact we can make the following definition.

Definition 3.2. Let K be an augmented directed complex with a basis and let b be a basis element. Then the *atom* associated to b is the element $\langle b \rangle$ of μK such that $\langle b \rangle_n^\alpha = 0$ for $n > |b|$, such that $\langle b \rangle_{|b|}^\alpha = b$, and such that $\langle b \rangle_n^- \wedge \langle b \rangle_n^+ = 0$ for $n < |b|$. The *dimension* of the atom $\langle b \rangle$ is the dimension of the corresponding basis element b .

From Proposition 2.7 we deduce the following result.

Proposition 3.3. Let K be an augmented directed complex with a basis. Then the atoms form a basis for the abelian group μK , the n -dimensional atoms form a basis for $(\mu K)_n / (\mu K)_{n-1}$, and the atoms of dimension greater than n form a basis for $\mu K / (\mu K)_n$.

If $\langle b \rangle$ is an atom, then $\langle b \rangle_n^\alpha \in K_n^*$ for all n by construction. An atom $\langle b \rangle$ is therefore in νK if and only if $\varepsilon \langle b \rangle_0^- = \varepsilon \langle b \rangle_0^+ = 1$. This leads us to the following definition.

Definition 3.4. A basis B for an augmented directed complex is *unital* if $\varepsilon \langle b \rangle_0^- = \varepsilon \langle b \rangle_0^+ = 1$ for every $b \in B$.

For the equivalence theorem of Section 5 we need bases which are unital and are also loop-free in the sense of the following definition.

Definition 3.5. A basis B for an augmented directed complex is *loop-free* if there are partial orderings \leq_0, \leq_1, \dots on B such that $a <_n b$ whenever $\langle a \rangle_n^+ \wedge \langle b \rangle_n^- > 0$ and $|a|, |b| > n$.

Note that the partial orderings \leq_n are quite different from the partial orderings on the individual chain groups.

In practice, one usually has a stronger condition.

Definition 3.6. A basis B for an augmented directed complex is *strongly loop-free* if there is a partial ordering $\leq_{\mathbb{N}}$ on B such that $a <_{\mathbb{N}} b$ whenever $a \leq \partial^- b$ or $\partial^+ a \geq b$.

The two notions of loop-freeness are related as follows.

Proposition 3.7. If a basis for an augmented directed complex is strongly loop-free, then it is loop-free.

Proof. Let $\leq_{\mathbf{N}}$ be a partial ordering with the property required for strong loop-freeness. For each n , we will show that $\leq_{\mathbf{N}}$ has the property required for \leq_n in the definition of loop-freeness. In other words, we suppose that $\langle a \rangle_n^+ \wedge \langle b \rangle_n^- > 0$ with $|a|, |b| > n$ for some n , and we show that $a <_{\mathbf{N}} b$. Indeed we can choose a basis element c with $\langle a \rangle_n^+ \geq c$ and $c \leq \langle b \rangle_n^-$, and we will show that $a <_{\mathbf{N}} c <_{\mathbf{N}} b$.

To show that $c <_{\mathbf{N}} b$, observe that, by the construction of $\langle b \rangle$, we must have $c \leq \partial^- c'$ for some basis element c' with $c' \leq \langle b \rangle_{n+1}^-$, and we then have $c <_{\mathbf{N}} c'$. If $n + 1 < |b|$ then we repeat this argument, and eventually we get $c <_{\mathbf{N}} \dots <_{\mathbf{N}} c''$ with $c'' \leq \langle b \rangle_{|b|}^-$. But $\langle b \rangle_{|b|}^- = b$, so $c'' = b$, and we have got $c <_{\mathbf{N}} b$ as claimed. The proof that $a <_{\mathbf{N}} c$ is similar. \square

Example 3.8. Let K be the chain complex of a simplicial set. Then K is an augmented chain complex with a distinguished basis, so it can be regarded as an augmented directed complex with a basis. In particular, let $\Delta[p]$ be the chain complex of the standard p -simplex, so that $\Delta[p]$ has the following structure: the basis elements are the ordered $(n + 1)$ -tuples of integers (v_0, \dots, v_n) with $0 \leq v_0 < v_1 < \dots < v_n \leq p$; the dimension of (v_0, \dots, v_n) is n ; the boundary $\partial: \Delta[p]_n \rightarrow \Delta[p]_{n-1}$ is given by

$$\partial = \partial_0 - \partial_1 + \partial_2 - \dots + (-1)^n \partial_n,$$

where

$$\partial_i(v_0, \dots, v_n) = (v_0, \dots, v_{i-1}, v_{i+1}, \dots, v_n);$$

the augmentation is given by $\varepsilon(v_0) = 1$. For $m \leq n$ one finds that

$$\langle (v_0, \dots, v_n) \rangle_m^\alpha = \sum \partial_{i(1)} \dots \partial_{i(n-m)}(v_0, \dots, v_n),$$

where the sum runs over $(n - m)$ -tuples such that $0 \leq i(1) < \dots < i(n - m) \leq n$ and the parities $(-)^{i(1)}, \dots, (-)^{i(n-m)}$ form the alternating sequence $\alpha, -\alpha, \alpha, \dots$. We have therefore recovered Street's oriented simplexes [11]. The basis is unital, because $\langle (v_0, \dots, v_n) \rangle_0^- = (v_0)$ and $\langle (v_0, \dots, v_n) \rangle_0^+ = (v_n)$. The basis is also strongly loop-free under the total ordering given recursively as follows: $(v_0, \dots, v_n) <_{\mathbf{N}} (w_0, \dots, w_m)$ if

$$v_0 < w_0,$$

or if

$$v_0 = w_0 \text{ and } n = 0 \text{ and } m > 0,$$

or if

$$v_0 = w_0 \text{ and } n > 0 \text{ and } m > 0 \text{ and } (v_1, \dots, v_n) >_{\mathbf{N}} (w_1, \dots, w_m).$$

Example 3.9. Let B be a finite non-empty totally ordered set in which each element b is assigned a nonnegative integer dimension $|b|$. Suppose also that the initial and final elements have dimension 0 and that adjacent elements have dimensions differing by 1. For $b \in B$ with $|b| > 0$, let $\delta^- b$ be the last element of dimension $|b| - 1$ to come before b , and let $\delta^+ b$ be the first element of dimension $|b| - 1$ to come after b ; the hypotheses ensure that these elements exist. The hypotheses also ensure that the elements between $\delta^- b$ and $\delta^+ b$ have dimension at least $|b|$, and for

$|b| > 1$ it follows that $\delta^\alpha \delta^{-b} = \delta^\alpha \delta^{+b}$. By taking $\partial b = \delta^{+b} - \delta^{-b}$ for $|b| > 0$ and $\varepsilon b = 1$ for $|b| = 0$ we define an augmented directed complex with basis B . Clearly $\langle b \rangle_n^\alpha$ is a basis element for $n \leq |b|$, so the basis is unital. It is also strongly loop-free under the original total ordering.

In particular, let p be a nonnegative integer and let the sequence of dimensions of the elements of B be

$$0, 1, \dots, p-1, p, p-1, \dots, 1, 0.$$

Then the augmented directed complex is called the p -dimensional globe and denoted $G[p]$. Let x denote the p -dimensional basis element; then $d_p^- \langle x \rangle = d_p^+ \langle x \rangle = \langle x \rangle$, and the atoms other than $\langle x \rangle$ are the elements $d_i^\alpha \langle x \rangle$ for $i < p$.

A p -dimensional globe is in a sense free on a p -dimensional generator. In a similar way we can get an augmented directed complex $G[p; n]$ free on a $\#_n$ -composable pair of p -dimensional elements: we take the sequence of dimensions of the elements of B to run from 0 up to p , then down to $\min\{p, n\}$, then up to p , then down to 0. Let the p -dimensional basis elements in order be x and y (they coincide if $p \leq n$); then $d_p^- \langle x \rangle = d_p^+ \langle x \rangle = \langle x \rangle$, $d_p^- \langle y \rangle = d_p^+ \langle y \rangle = \langle y \rangle$, and $d_n^+ \langle x \rangle = d_n^- \langle y \rangle$. The atoms other than $\langle x \rangle$ and $\langle y \rangle$ are the $d_i^\alpha \langle x \rangle$ and $d_i^\alpha \langle y \rangle$ for $i < p$, with the identifications $d_i^\alpha \langle x \rangle = d_i^\alpha \langle y \rangle$ for $i < \min\{p, n\}$ and with the identification $d_n^+ \langle x \rangle = d_n^- \langle y \rangle$ if $n < p$.

There is a similar augmented directed complex $G[p; n, n]$ free on a $\#_n$ -composable triple of p -dimensional elements: the sequence of dimensions is now from 0 up to p , down to $\min\{p, n\}$, up to p , down to $\min\{p, n\}$, up to p , down to 0. For $m < n$ there is also an augmented directed complex $G[p; n, m, n]$ free on a composable quadruple of p -dimensional elements in the configuration $(x \#_n y) \#_m (x' \#_n y')$ of Definition 2.1(vii): the sequence of dimensions is now from 0 up to p , down to $\min\{p, n\}$, up to p , down to $\min\{p, m\}$, up to p , down to $\min\{p, n\}$, up to p , down to 0.

Example 3.10. The category of augmented chain complexes has a symmetric monoidal structure under the tensor product $(K, L) \mapsto K \otimes L$, where

$$\begin{aligned} (K \otimes L)_n &= \bigoplus_i K_i \otimes L_{n-i}, \\ \partial(x \otimes y) &= \partial x \otimes y + (-1)^{|x|} x \otimes \partial y, \\ \varepsilon(x \otimes y) &= (\varepsilon x)(\varepsilon y) \text{ for } |x| = |y| = 0 \end{aligned}$$

(we write $|x| = i$ if $x \in K_i$, etc.). The identity object is the 0-dimensional globe $G[0]$ of Example 3.9. We extend this structure to a monoidal structure on augmented directed complexes as follows: $(K \otimes L)_n^*$ is the submonoid of $(K \otimes L)_n$ generated by the elements $x \otimes y$ with $x \in K_i^*$ and $y \in L_{n-i}^*$. Note that the monoidal structure on augmented directed complexes is not symmetric: the standard switch morphism $x \otimes y \mapsto (-1)^{|x||y|} (y \otimes x)$ does not always map $(K \otimes L)_n^*$ into $(L \otimes K)_n^*$.

Suppose now that K and L are augmented directed complexes with bases A and B . Then $K \otimes L$ has a basis C consisting of the elements $a \otimes b$ for $a \in A$ and $b \in B$.

One finds that

$$\langle a \otimes b \rangle_n^\alpha = \sum_{i=0}^n \langle a \rangle_i^\alpha \otimes \langle b \rangle_{n-i}^{(-)^i \alpha};$$

in particular $\langle a \otimes b \rangle_0^\alpha = \langle a \rangle_0^\alpha \otimes \langle b \rangle_0^\alpha$. If A and B are unital, then it follows that C is unital. Similarly, if A and B are strongly loop-free under partial orderings $\leq_{\mathbf{N}}$, then C is strongly loop-free under the partial ordering $\leq_{\mathbf{N}}$ such that $a \otimes b \leq_{\mathbf{N}} a' \otimes b'$ for

$$a <_{\mathbf{N}} a',$$

or for

$$a = a' \text{ and } |a| \text{ even and } b \leq_{\mathbf{N}} b',$$

or for

$$a = a' \text{ and } |a| \text{ odd and } b \geq_{\mathbf{N}} b'.$$

In particular, for $p \geq 0$, let $Q[p]$ be the chain complex of the p -dimensional cube. Then $Q[p]$ is the p -fold tensor power of the one-dimensional globe $G[1]$, so $Q[p]$ is an augmented directed complex with a strongly loop-free unital basis.

4. Bases for ω -categories

We will now describe bases for ω -categories, corresponding to bases for augmented directed complexes. The required properties can be described directly in terms of ω -categories, but it is often easier to work in the augmented directed complexes got by applying λ .

We begin with a particular kind of generating set, analogous to a spanning set. Recall that if C is an ω -category then C_n is the sub- ω -category $d_n^- C = d_n^+ C$.

Definition 4.1. An ω -category C is *composition-generated* by a subset E if each member e of E is assigned a nonnegative integer dimension $|e|$ and if for $n \geq 0$ the sub- ω -category C_n is generated under the composition operations $\#_m$ by the elements of E of dimension at most n .

Composition-generation should be distinguished from ordinary generation, where one uses the operations d_m^α as well as the composition operations.

Note that if e is a composition-generator for an ω -category C then $e \in C_{|e|}$, so that $d_n^\alpha e = e$ for $n \geq |e|$.

There are standard forms for the elements of ω -categories with composition-generators, as follows.

Proposition 4.2. *Let C be an ω -category with a set of composition-generators and let x be a member of C . Then x is a generator or x has an expression $x = x_1 \#_r \dots \#_r x_k$ with $r \geq 0$ and $k \geq 2$ such that the x_i are composites of generators, each x_i has exactly one factor of dimension greater than r , and at most one of the x_i has a factor of dimension greater than $r + 1$.*

Proof. It suffices to prove the result when x is a composite, say $x = y \#_n z$, in which y and z have expressions of the required form. If the expression for y involves no

factors of dimension greater than n , then y is an identity for $\#_n$ by Definition 2.1(v), so $x = z$ and the result therefore holds for x . A similar argument applies if the expression for z involves no factors of dimension greater than n .

From now on, assume that the expressions for y and z both involve factors of dimension greater than n . Let r be the largest integer such that the expressions for y and z together involve more than one factor of dimension greater than r ; thus $r \geq n$. Suppose that the expressions for y and z have k and l factors respectively of dimension greater than r ; thus $k + l \geq 2$. Then there is a decomposition

$$y = y_1 \#_r \dots \#_r y_k \#_r d_r^+ y \#_r \dots \#_r d_r^+ y,$$

with l appearances of $d_r^+ y$, such that y_i has exactly one factor of dimension greater than r (if $k = 0$ then $y = d_r^+ y$, so this holds trivially; if $k = 1$ we take $y_1 = y$; if $k > 1$ then $y_1 \#_r \dots \#_r y_k$ is the given expression for y). Similarly there is a decomposition

$$z = d_r^- z \#_r \dots \#_r d_r^- z \#_r z_1 \#_r \dots \#_r z_l,$$

with k factors equal to $d_r^- z$, such that z_j has exactly one factor of dimension greater than r . The choice of r ensures that the y_i and z_j have at most one factor of dimension $r + 1$ between them. If now $n = r$ then the desired decomposition of $x = y \#_r z$ is given by

$$x = y_1 \#_r \dots \#_r y_k \#_r z_1 \#_r \dots \#_r z_l;$$

if $n < r$ then Definition 2.1(vii) gives

$$\begin{aligned} x &= (y_1 \#_r \dots \#_r y_k \#_r d_r^+ y \#_r \dots \#_r d_r^+ y) \#_n (d_r^- z \#_r \dots \#_r d_r^- z \#_r z_1 \#_r \dots \#_r z_l) \\ &= (y_1 \#_n d_r^- z) \#_r \dots \#_r (y_k \#_n d_r^- z) \#_r (d_r^+ y \#_n z_1) \#_r \dots \#_r (d_r^+ y \#_n z_l), \end{aligned}$$

which is a decomposition of the required form because $d_r^- z$ and $d_r^+ y$ are composites of generators of dimension at most r .

This completes the proof. □

Composition-generators for an ω -category C produce generators for $(\lambda C)_n$ and $(\lambda C)_n^*$ as follows.

Proposition 4.3. *Let C be an ω -category with a set of composition-generators E . Then the abelian group $(\lambda C)_n$ and the submonoid $(\lambda C)_n^*$ are generated by the elements $[e]_n$ for e an n -dimensional element of E .*

Proof. The abelian group $(\lambda C)_n$ and the submonoid $(\lambda C)_n^*$ are generated by the elements $[x]_n$ for $x \in C_n$, and the elements of C_n are the composites of the members of E of dimension at most n . If $m \geq n$ then the elements of C_n are identities for $\#_m$, because $d_m^\beta d_n^\alpha = d_n^\alpha$; it therefore suffices to use the composition operators $\#_m$ for $m < n$. These operations become addition in $(\lambda C)_n$, so $(\lambda C)_n$ and $(\lambda C)_n^*$ are generated by the elements $[e]_n$ for $e \in E$ with $|e| \leq n$. Finally $[e]_n = 0$ for $|e| < n$ by Proposition 2.5; therefore $(\lambda C)_n$ and $(\lambda C)_n^*$ are generated by the elements $[e]_n$ for $e \in E$ with $|e| = n$. □

We now define a basis for an ω -category C in terms of a basis for the augmented directed complex λC .

Definition 4.4. A *basis* for an ω -category C is a set of composition-generators E such λC has a basis and the function $e \mapsto [e]_{|e|}$ maps E bijectively onto the basis for λC .

Because of Proposition 4.3, a composition-generating set E is a basis if and only if the elements $[e]_{|e|}$ for $e \in E$ are distinct and linearly independent.

Let C be an ω -category with a basis E . Then λC has a basis B , and we can use properties of B to define properties of E .

Definition 4.5. Let E be a basis for an ω -category C . Then

- (i) E is *atomic* if $[d_n^- e]_n \wedge [d_n^+ e]_n = 0$ for $e \in E$ and $n < |e|$;
- (ii) E is *loop-free* if the basis for λC is loop-free;
- (iii) E is *strongly loop-free* if the basis for λC is strongly loop-free.

The point of atomicity is as follows.

Proposition 4.6. Let E be an atomic basis for an ω -category C . Then $\eta e = \langle [e]_{|e|} \rangle$ for each e in E , and the basis for λC is unital.

Proof. Recall from Definition 2.9 that $(\eta e)_n^\alpha = [d_n^\alpha e]_n$ for all n and α . If $n > |e|$ then $[d_n^\alpha e]_n = [e]_n = 0$ by Proposition 2.5. If $n = |e|$ then $[d_n^\alpha e]_n = [e]_{|e|}$. By comparing Definition 4.5(i) with Definition 3.2, we now see that $\eta e = \langle [e]_{|e|} \rangle$. Since $\eta e \in \nu \lambda C$, it follows that $\varepsilon \langle [e]_{|e|} \rangle_0^\alpha = \varepsilon (\eta e)_0^\alpha = 1$; therefore the basis for λC is unital. This completes the proof. \square

Example 4.7. Let p be a nonnegative integer, and let $F[p]$ be the ω -category with the following presentation: there is a single generator u and there are relations $d_p^- u = d_p^+ u = u$. The ω -categories $F[p]$ represent the elements of ω -categories, in the sense that there are natural bijections between $\text{hom}(F[p], C)$ and C_p . We have $d_i^\alpha u = u$ for $i \geq p$. Using Definition 2.1(iii) and (v), we see that $F[p]$ is composition-generated by u and the elements $d_i^\alpha u$ for $i < p$.

Now let $G[p]$ be the p -dimensional globe with p -dimensional basis element x as in Example 3.9. The atom $\langle x \rangle$ in $\nu G[p]$ satisfies the relations $d_p^- \langle x \rangle = d_p^+ \langle x \rangle = \langle x \rangle$, so there is a morphism $F[p] \rightarrow \nu G[p]$ given by $u \mapsto \langle x \rangle$. The adjoint $\lambda F[p] \rightarrow G[p]$ maps the generators $[u]_p$ and $[d_i^\alpha u]_i$ for $\lambda F[p]$ (see Proposition 4.3) bijectively to the basis elements for $G[p]$, and it follows that the morphism $\lambda F[p] \rightarrow G[p]$ is an isomorphism. It then follows that the elements u and $d_i^\alpha u$ form a basis for $F[p]$, and one can check that this basis is atomic. It is also strongly loop-free, because the basis for $G[p]$ is strongly loop-free.

There are similar ω -categories $F[p; n]$, $F[p; n, n]$ and $F[p; n, m, n]$ (where $m < n$) corresponding to the other augmented directed complexes of Example 3.9. These ω -categories all have strongly loop-free atomic bases. There is a presentation for $F[p; n]$ given by

$$\langle u, v : d_p^\alpha u = u, d_p^\alpha v = v, d_n^+ u = d_n^- v \rangle,$$

there is a presentation for $F[p; n, n]$ given by

$$\langle u, v, w : d_p^\alpha u = u, d_p^\alpha v = v, d_p^\alpha w = w, d_n^+ u = d_n^- v, d_n^+ v = d_n^- w \rangle,$$

and there is a presentation for $F[p; n, m, n]$ given by

$$\langle u, v, u', v' : d_p^\alpha u = u, d_p^\alpha v = v, d_p^\alpha u' = u', d_p^\alpha v' = v', \\ d_n^+ u = d_n^- v, d_m^+ v = d_m^- u', d_n^+ u' = d_n^- v' \rangle.$$

5. The adjoint equivalence

From Definition 4.4, Definition 4.5 and Proposition 4.6, if an ω -category C has a loop-free atomic basis then λC has a loop-free unital basis. We will now show that ω -categories with loop-free atomic bases and augmented directed complexes with loop-free unital bases are equivalent under the adjoint functors λ and ν .

We begin by considering decompositions in νK , where K is an augmented directed complex with a basis. The basic existence result is as follows.

Proposition 5.1. *Let K be an augmented directed complex with a basis and let x be a member of νK which is congruent to a non-trivial sum of atoms modulo $(\mu K)_r$ for some $r \geq 0$, say*

$$x \equiv \langle b_1 \rangle + \dots + \langle b_k \rangle \pmod{(\mu K)_r}$$

with $k \geq 1$. If $\langle b_i \rangle_r^+ \wedge \langle b_j \rangle_r^- = 0$ for $i > j$, then there is a decomposition

$$x = x_1 \#_r \dots \#_r x_k$$

with $x_i \in \nu K$ such that $x_i \equiv \langle b_i \rangle \pmod{(\mu K)_r}$.

Proof. The congruence gives

$$x = \langle b_1 \rangle + \dots + \langle b_k \rangle + z$$

for some $z \in (\mu K)_r$; thus $d_r^\alpha z = z$. For $1 \leq i \leq k$ let x_i be the element of μK given by

$$x_i = d_r^+[\langle b_1 \rangle + \dots + \langle b_{i-1} \rangle] + \langle b_i \rangle + d_r^-[\langle b_{i+1} \rangle + \dots + \langle b_k \rangle] + z,$$

and for $1 \leq i \leq k-1$ let

$$y_i = d_r^+[\langle b_1 \rangle + \dots + \langle b_i \rangle] + d_r^-[\langle b_{i+1} \rangle + \dots + \langle b_k \rangle] + z.$$

Then $d_r^+ x_i = y_i = d_r^- x_{i+1}$ because $d_r^\beta d_r^\alpha = d_r^\alpha$, and we also have

$$x = x_1 - y_1 + x_2 - y_2 + \dots + x_k,$$

so there is a decomposition

$$x = x_1 \#_r \dots \#_r x_k$$

in μK . It is clear that $x_i \equiv \langle b_i \rangle \pmod{(\mu K)_r}$, and it remains to show that $x_i \in \nu K$. We must therefore show that $\varepsilon(x_i)_0^\alpha = 1$ and that $(x_i)_n^\alpha \geq 0$.

As to the augmentation, we have $\varepsilon(d_r^\beta \langle b_j \rangle)_0^\alpha = \varepsilon \langle b_j \rangle_0^\alpha$ for all j and β , so $\varepsilon(x_i)_0^\alpha = \varepsilon x_0^\alpha = 1$.

As to the $(x_i)_n^\alpha$, if $n < r$ then $(x_i)_n^\alpha = x_n^\alpha \geq 0$, and if $n > r$ then $(x_i)_n^\alpha = \langle b_i \rangle_n^\alpha \geq 0$; it therefore remains to consider the case $n = r$. Now $(x_1)_r^- = x_r^- \geq 0$ and

$(x_k)_r^+ = x_r^+ \geq 0$, and for $1 \leq i \leq k-1$ we have

$$\begin{aligned} (x_i)_r^+ &= (x_{i+1})_r^- = x_r^- + [\langle b_1 \rangle_r^+ + \dots + \langle b_i \rangle_r^+] - [\langle b_1 \rangle_r^- + \dots + \langle b_i \rangle_r^-] \\ &= x_r^+ + [\langle b_{i+1} \rangle_r^- + \dots + \langle b_k \rangle_r^-] - [\langle b_{i+1} \rangle_r^+ + \dots + \langle b_k \rangle_r^+]. \end{aligned}$$

But $x_r^\alpha \geq 0$, $\langle b_j \rangle_r^\alpha \geq 0$, and

$$[\langle b_1 \rangle_r^- + \dots + \langle b_i \rangle_r^-] \wedge [\langle b_{i+1} \rangle_r^+ + \dots + \langle b_k \rangle_r^+] = 0,$$

so $(x_i)_r^\alpha \geq 0$ in all cases. This completes the proof. \square

To find situations in which Proposition 5.1 can be applied, we use the following result.

Proposition 5.2. *Let K be an augmented directed complex with a basis and let x be a member of νK .*

(i) *If $x \equiv 0 \pmod{(\mu K)_{r+1}}$ then*

$$x \equiv \langle c_1 \rangle + \dots + \langle c_l \rangle \pmod{(\mu K)_r}$$

for some $(r+1)$ -dimensional atoms $\langle c_1 \rangle, \dots, \langle c_l \rangle$.

(ii) *If $x \equiv m\langle a \rangle \pmod{(\mu K)_{r+1}}$ for some positive integer m and for some atom $\langle a \rangle$ with $|a| > r+1$, then*

$$x \equiv m\langle a \rangle + \langle c_1 \rangle + \dots + \langle c_l \rangle \pmod{(\mu K)_r}$$

for some $(r+1)$ -dimensional atoms $\langle c_1 \rangle, \dots, \langle c_l \rangle$.

(iii) *If $x = y \#_r z$ for some $y, z \in \nu K$ such that $y \equiv \langle a \rangle$ and $z \equiv \langle b \rangle \pmod{(\mu K)_r}$ with $|a|, |b| > r$ and $\langle a \rangle_r^+ \wedge \langle b \rangle_r^- = 0$, then*

$$x \equiv \langle a \rangle + \langle b \rangle + \langle c_1 \rangle + \dots + \langle c_l \rangle \pmod{(\mu K)_{r-1}}$$

for some r -dimensional atoms $\langle c_1 \rangle, \dots, \langle c_l \rangle$.

Proof. (i) Since $x \in (\mu K)_{r+1}$, we have $x_{r+1}^- = x_{r+1}^+ = w$ for some $w \in K_{r+1}$. Since $x \in \nu K$, we have $w \geq 0$, so that w is a sum of $(r+1)$ -dimensional basis elements $c_1 + \dots + c_l$. It now follows that $x \equiv \langle c_1 \rangle + \dots + \langle c_l \rangle \pmod{(\mu K)_r}$, as required.

(ii) Here $x - m\langle a \rangle$ is in $(\mu K)_{r+1}$. As in the proof of part (i), we get $x_{r+1}^- - m\langle a \rangle_{r+1}^- = x_{r+1}^+ - m\langle a \rangle_{r+1}^+ = w$ for some $w \in K_{r+1}$, and it suffices to show that $w \geq 0$. But this holds because $x_{r+1}^- \geq 0$, $x_{r+1}^+ \geq 0$ and $\langle a \rangle_{r+1}^- \wedge \langle a \rangle_{r+1}^+ = 0$.

(iii) Here

$$x_r^- - \langle a \rangle_r^- - \langle b \rangle_r^- = x_r^+ - \langle a \rangle_r^+ - \langle b \rangle_r^+ = z_r^- - \langle a \rangle_r^+ - \langle b \rangle_r^- = w$$

for some $w \in K_r$, since $x \equiv y + z \equiv \langle a \rangle + \langle b \rangle \pmod{(\mu K)_r}$, since $x_r^+ = z_r^+$, and since $z - \langle b \rangle \in (\mu K)_r$. As before, it suffices to show that $w \geq 0$. But this holds because $x_r^- \geq 0$, $x_r^+ \geq 0$, $z_r^- \geq 0$ and

$$\langle a \rangle_r^- \wedge \langle a \rangle_r^+ = \langle b \rangle_r^- \wedge \langle b \rangle_r^+ = \langle a \rangle_r^+ \wedge \langle b \rangle_r^- = 0.$$

\square

To construct decompositions we use Proposition 5.2(i) together with the case $m = 1$ of Proposition 5.2(ii). It is convenient to make the following definition.

Definition 5.3. Let K be an augmented directed complex with a basis, and let x be a member of νK . Then the *decomposition index* of x is the smallest integer $r \geq -1$ such that x is congruent to zero or an atom modulo $(\mu K)_{r+1}$.

In Definition 5.3, note that the decomposition index exists because $x \equiv 0 \pmod{(\mu K)_n}$ for all sufficiently large n .

By combining Propositions 5.1 and 5.2 we get the following result, analogous to part of Proposition 4.2.

Proposition 5.4. *Let K be an augmented directed complex with a loop-free basis, and let x be a member of $(\nu K)_n$ with decomposition index $r \geq 0$. Then there is a decomposition*

$$x = x_1 \#_r \dots \#_r x_k$$

with $x_i \in (\nu K)_n$ and $x_i \equiv \langle b_i \rangle \pmod{(\mu K)_r}$, where $\langle b_1 \rangle, \dots, \langle b_k \rangle$ is a list of atoms such that $k \geq 2$, such that $|b_i| > r$ for all i , such that $|b_i| > r + 1$ for at most one value of i , and such that $\langle b_i \rangle_r^+ \wedge \langle b_j \rangle_r^- = 0$ for $i > j$.

Proof. We have x congruent to zero or an atom modulo $(\mu K)_{r+1}$. By Proposition 5.2(i) or (ii) there is a congruence

$$x \equiv \langle b_1 \rangle + \dots + \langle b_k \rangle \pmod{(\mu K)_r}$$

for some list of atoms $\langle b_i \rangle$ with $|b_i| > r$ for each i and with $|b_i| > r + 1$ for at most one value of i . Since the decomposition index is r , we have $k \geq 2$. Since the basis is loop-free we can assume the list ordered so that $\langle b_i \rangle_r^+ \wedge \langle b_j \rangle_r^- = 0$ for $i > j$. By Proposition 5.1 there is a decomposition $x = x_1 \#_r \dots \#_r x_k$ with $x_i \in \nu K$ and $x_i \equiv \langle b_i \rangle \pmod{(\mu K)_r}$. Finally, $|b_i| \leq n$ for each i , because $x \in (\nu K)_n$, and it follows that $x_i \in (\nu K)_n$. This completes the proof. \square

To complete the analogy with Proposition 4.2 we give the following result.

Proposition 5.5. *Let K be an augmented directed complex with a unital basis, and let x be a member of $(\nu K)_n$ with decomposition index -1 . Then x is an atom of dimension at most n .*

Proof. We have x congruent to zero or an atom modulo $(\mu K)_0$. By Proposition 5.2(i) or (ii) there is a congruence

$$x \equiv \langle b_1 \rangle + \dots + \langle b_k \rangle \pmod{(\mu K)_{-1}}$$

for some atoms $\langle b_i \rangle$. Since $(\mu K)_{-1} = 0$, this congruence is an equality, and it follows that

$$\varepsilon x_0^\alpha = \varepsilon \langle b_1 \rangle_0^\alpha + \dots + \varepsilon \langle b_k \rangle_0^\alpha.$$

But $\varepsilon x_0^\alpha = 1$ because $x \in \nu K$, and $\varepsilon \langle b_i \rangle_0^\alpha = 1$ for each i because the basis is unital, so $k = 1$. This means that x is equal to the atom $\langle b_1 \rangle$. The dimension of $\langle b_1 \rangle$ is at most n because $x \in (\nu K)_n$. \square

We can now prove one half of the equivalence.

Theorem 5.6. *Let K be an augmented directed complex with a loop-free unital basis. Then νK has a loop-free atomic basis consisting of its atoms, and the counit $\pi: \lambda\nu K \rightarrow K$ is an isomorphism.*

Proof. First we show that νK is composition-generated by its atoms. Indeed the atoms of dimension at most n lie in $(\nu K)_n$ because the basis is unital, so the composites of atoms of dimension at most n lie in $(\nu K)_n$. Conversely, let x be a member of $(\nu K)_n$ of decomposition index r . If $r \geq 0$, then x is a composite of members of $(\nu K)_n$ with lower decomposition index by Proposition 5.4, and if $r = -1$ then x is an atom of dimension at most n by Proposition 5.5. It follows by induction on r that x is a composite of atoms of dimension at most n . Therefore νK is composition-generated by its atoms.

Next we show that $\pi: \lambda\nu K \rightarrow K$ is an isomorphism. For each $n \geq 0$ we must show that $\pi: (\lambda\nu K)_n \rightarrow K_n$ is an isomorphism with $\pi[(\lambda\nu K)_n^*] = K_n^*$. Now it follows from Proposition 4.3 that $(\lambda\nu K)_n$ and $(\lambda\nu K)_n^*$ are generated as abelian group and submonoid by the elements $[\langle b \rangle]_n$ for $|b| = n$. The images of these elements under π are the n -dimensional basis elements, which generate K_n and K_n^* as abelian group and submonoid freely. It follows that $\pi: (\lambda\nu K)_n \rightarrow K_n$ is an isomorphism with $\pi[(\lambda\nu K)_n^*] = K_n^*$ as required.

Finally we show that the atoms form a basis for νK with the desired properties by verifying the conditions of Definitions 4.4 and 4.5. We have already shown that the atoms composition-generate νK . The elements $[\langle b \rangle]_{|b|}$ form a basis for $\lambda\nu K$, because their images under the isomorphism $\pi: \lambda\nu K \rightarrow K$ form a basis for K , so the function $\langle b \rangle \mapsto [\langle b \rangle]_{|b|}$ is a bijection from the set of atoms to a basis for $\lambda\nu K$. The atoms therefore form a basis for νK . We have $[d_n^-(b)]_n \wedge [d_n^+(b)]_n = 0$ for $n < |b|$ because

$$\pi[d_n^-(b)]_n \wedge \pi[d_n^+(b)]_n = \langle b \rangle_n^- \wedge \langle b \rangle_n^+ = 0;$$

therefore the basis for νK is atomic. The basis for $\lambda\nu K$ is loop-free because $\lambda\nu K \cong K$; therefore the basis for νK is loop-free. This completes the proof. \square

Conversely, if C is an ω -category with a loop-free atomic basis, then we want the unit $\eta: C \rightarrow \nu\lambda C$ to be an isomorphism. More generally, we will prove the following result.

Theorem 5.7. *Let C be an ω -category composition-generated by a set E , let K be an augmented directed complex with a loop-free unital basis, and let $\theta: C \rightarrow \nu K$ be a morphism which restricts to a dimension-preserving bijection from E to the atoms of νK . Then θ is an isomorphism.*

In a sense, Theorem 5.7 says that νK is freely composition-generated by its atoms. To prove it, we will show that the decompositions in Proposition 5.4 are as nearly as possible uniquely determined. We begin with the following observation.

Proposition 5.8. *Let K be an augmented directed complex. If*

$$x_1 \#_r \dots \#_r x_k = y_1 \#_r \dots \#_r y_k$$

in μK and if $x_i \equiv y_i \pmod{(\mu K)_r}$ for $1 \leq i \leq k$, then $x_i = y_i$ for $1 \leq i \leq k$.

Proof. We have

$$(x_1 - y_1) \#_r \dots \#_r (x_k - y_k) = 0$$

and $x_1 - y_1 \in (\mu K)_r$, so

$$x_1 - y_1 = d_r^-(x_1 - y_1) = d_r^-(0) = 0,$$

and it follows that $x_1 = y_1$. We then have $d_r^-(x_2 - y_2) = d_r^+(x_1 - y_1) = 0$, so $x_2 = y_2$ by a similar argument, and so on. \square

Because of Proposition 5.8, the decomposition in Proposition 5.4 is uniquely determined by the ordered list of atoms $\langle b_1 \rangle, \dots, \langle b_k \rangle$. Up to permutation this list is determined by x , because $x \equiv \langle b_1 \rangle + \dots + \langle b_k \rangle \pmod{(\mu K)_r}$; however, it may be possible to reorder the list.

In particular, suppose that $k = 2$, so that $x \equiv \langle b_1 \rangle + \langle b_2 \rangle \pmod{(\mu K)_r}$ with $\min\{|b_1|, |b_2|\} = r + 1$. Suppose further that $r > 0$, and that x has a $\#_{r-1}$ -decomposition $x = x_1^* \#_{r-1} x_2^*$ with $x_1^* \equiv \langle b_1 \rangle$ and $x_2^* \equiv \langle b_2 \rangle \pmod{(\mu K)_r}$. Then it follows from Definition 2.1(iv) and (vii) that there are $\#_r$ -decompositions

$$x = (x_1^* \#_{r-1} d_r^- x_2^*) \#_r (d_r^+ x_1^* \#_{r-1} x_2^*) = (d_r^- x_1^* \#_{r-1} x_2^*) \#_r (x_1^* \#_{r-1} d_r^+ x_2^*)$$

with $(x_1^* \#_{r-1} d_r^- x_2^*) \equiv \langle b_1 \rangle$ and $(d_r^+ x_1^* \#_{r-1} x_2^*) \equiv \langle b_2 \rangle \pmod{(\mu K)_r}$. For elements of this type, one can therefore transpose the atoms. We will now show that any two consecutive atoms in the ‘wrong’ order can be transposed in this way.

Proposition 5.9. *Let K be an augmented directed complex with a loop-free unital basis. Suppose that $x = y \#_r z$ is a composite in νK with $y \equiv \langle a \rangle$ and $z \equiv \langle b \rangle \pmod{(\mu K)_r}$, where $\langle a \rangle$ and $\langle b \rangle$ are atoms such that $|a|, |b| > r$ and $\langle a \rangle_r^+ \wedge \langle b \rangle_r^- = 0$. Then $r > 0$ and there is a decomposition $x = y^* \#_{r-1} z^*$ or $x = z^* \#_{r-1} y^*$ in νK with $y^* \equiv \langle a \rangle$ and $z^* \equiv \langle b \rangle \pmod{(\mu K)_r}$.*

Proof. By Proposition 5.2(iii),

$$x \equiv \langle a \rangle + \langle b \rangle + \langle c_1 \rangle + \dots + \langle c_l \rangle \pmod{(\mu K)_{r-1}}$$

for some r -dimensional atoms $\langle c_i \rangle$. If $r = 0$ then this congruence is an equality and we get $\varepsilon x_0^\alpha = l + 2 > 1$, which is absurd. Therefore $r > 0$. Since the basis is loop-free we can order the list $\langle a \rangle, \langle b \rangle, \langle c_1 \rangle, \dots, \langle c_l \rangle$ so that Proposition 5.1 applies. We get a decomposition $x = x_1 \#_{r-1} \dots \#_{r-1} x_{l+2}$ in νK such that modulo $(\mu K)_{r-1}$ the factors are congruent to $\langle a \rangle, \langle b \rangle, \langle c_1 \rangle, \dots, \langle c_l \rangle$ in some order. Modulo $(\mu K)_r$ it follows that one of the factors is congruent to $\langle a \rangle$, another factor is congruent to $\langle b \rangle$, and the others are congruent to zero. By grouping the factors appropriately, we get a decomposition $x = y^* \#_{r-1} z^*$ or $x = z^* \#_{r-1} y^*$ of the required form. \square

In the proof of Theorem 5.7, we show that elements of νK have unique inverse images by induction on their decomposition indices. In the inductive step we use the following lemma.

Lemma 5.10. *Let K be an augmented directed complex with a loop-free unital basis, let C be an ω -category, let r be a nonnegative integer, and let $\theta: C \rightarrow \nu K$ be a morphism such that elements of νK with decomposition index less than r have*

unique inverse images in C . If $\xi = \eta \#_r \zeta$ is a composite in C such that $\theta(\eta) \equiv \langle a \rangle$ and $\theta(\zeta) \equiv \langle b \rangle$ modulo $(\mu K)_r$, where $\langle a \rangle$ and $\langle b \rangle$ are atoms such that $|a|, |b| > r$ and $\langle a \rangle_r^+ \wedge \langle b \rangle_r^- = 0$, then there is a decomposition $\xi = \zeta' \#_r \eta'$ in C such that $\theta(\zeta') \equiv \langle b \rangle$ and $\theta(\eta') \equiv \langle a \rangle$ modulo $(\mu K)_r$.

Proof. By Proposition 5.9, $r > 0$ and there is a decomposition $\theta(\xi) = y^* \#_{r-1} z^*$ or $\theta(\xi) = z^* \#_{r-1} y^*$ in νK with $y^* \equiv \langle a \rangle$ and $z^* \equiv \langle b \rangle$ modulo $(\mu K)_r$. For definiteness, suppose that $\theta(\xi) = y^* \#_{r-1} z^*$. Then y^* and z^* have decomposition indices less than r , so they have unique inverse images η^* and ζ^* . We now get

$$\theta(d_{r-1}^+ \eta^*) = d_{r-1}^+ y^* = d_{r-1}^- z^* = \theta(d_{r-1}^- \zeta^*),$$

and the element $d_{r-1}^+ y^* = d_{r-1}^- z^*$ clearly has decomposition index less than $r-1$, so $d_{r-1}^+ \eta^* = d_{r-1}^- \zeta^*$ by the uniqueness part of the hypothesis. It follows that η^* and ζ^* have a composite $\eta^* \#_{r-1} \zeta^*$. We now have $\theta(\xi) = \theta(\eta^* \#_{r-1} \zeta^*)$, and we claim that $\xi = \eta^* \#_{r-1} \zeta^*$. Indeed it follows from Definition 2.1(vii) that

$$\theta(\eta) \#_r \theta(\zeta) = \theta(\xi) = \theta(\eta^* \#_{r-1} \zeta^*) = \theta(\eta^* \#_{r-1} d_r^- \zeta^*) \#_r \theta(d_r^+ \eta^* \#_{r-1} \zeta^*)$$

with

$$\theta(\eta) \equiv \langle a \rangle \equiv y^* \equiv \theta(\eta^*) \equiv \theta(\eta^* \#_{r-1} d_r^- \zeta^*) \pmod{(\mu K)_r}$$

and

$$\theta(\zeta) \equiv \langle b \rangle \equiv z^* \equiv \theta(\zeta^*) \equiv \theta(d_r^+ \eta^* \#_{r-1} \zeta^*) \pmod{(\mu K)_r},$$

so $\theta(\eta) = \theta(\eta^* \#_{r-1} d_r^- \zeta^*)$ and $\theta(\zeta) = \theta(d_r^+ \eta^* \#_{r-1} \zeta^*)$ by Proposition 5.8. Clearly $\theta(\eta)$ and $\theta(\zeta)$ have decomposition indices less than r , so $\eta = \eta^* \#_{r-1} d_r^- \zeta^*$ and $\zeta = d_r^+ \eta^* \#_{r-1} \zeta^*$ by the uniqueness part of the hypothesis. Using Definition 2.1(vii) again, we get $\xi = \eta \#_r \zeta = \eta^* \#_{r-1} \zeta^*$ as claimed. It now follows that

$$\xi = (d_r^- \eta^* \#_{r-1} \zeta^*) \#_r (\eta^* \#_{r-1} d_r^+ \zeta^*) = \zeta' \#_r \eta',$$

say, with $\theta(\zeta') \equiv \theta(\zeta^*) \equiv \langle b \rangle$ and $\theta(\eta') \equiv \theta(\eta^*) \equiv \langle a \rangle$ modulo $(\mu K)_r$. This completes the proof. \square

Proof of Theorem 5.7. We show that θ is an isomorphism by showing that each element x of νK has a unique inverse image under θ . We use induction on the decomposition index of x .

Suppose first that x has decomposition index -1 . By Proposition 5.5, x is an atom. From Proposition 4.2, we see that the inverse images of x must be generators. Since θ maps the set of generators bijectively onto the set of atoms, it follows that x has a unique inverse image.

Now suppose that x has decomposition index $r \geq 0$. By Proposition 5.4 there is a decomposition

$$x = x_1 \#_r \dots \#_r x_k$$

with $k \geq 2$, with $x_i \in \nu K$, and with $x_i \equiv \langle b_i \rangle \pmod{(\mu K)_r}$, such that $|b_i| > r$ for all i , such that $|b_i| > r+1$ for at most one value of i , and such that $\langle b_i \rangle_r^+ \wedge \langle b_j \rangle_r^- = 0$ for $i > j$. The x_i clearly have decomposition indices less than r , so they have unique

inverse images ξ_i by the inductive hypothesis. As in the proof of Lemma 5.10,

$$\theta(d_r^+ \xi_{i-1}) = d_r^+ x_{i-1} = d_r^- x_i = \theta(d_r^- \xi_i)$$

and the decomposition index of $d_r^+ x_{i-1} = d_r^- x_i$ is less than r , so $d_r^+ \xi_{i-1} = d_r^- \xi_i$ by the uniqueness part of the inductive hypothesis, and it follows that there is a composite

$$\xi = \xi_1 \#_r \dots \#_r \xi_k.$$

Then ξ is an inverse image for x .

To complete the proof, let ξ' be any inverse image for x ; we must show that $\xi' = \xi$. We can express ξ' as in Proposition 4.2. From the form of x , this must mean that $\xi' = \xi'_1 \#_r \dots \#_r \xi'_k$ with the $\theta(\xi'_i)$ congruent to the $\langle b_i \rangle$ in some order modulo $(\mu K)_r$. By repeated application of Lemma 5.10, we can change back to the original order. This gives us a decomposition

$$\xi' = \xi''_1 \#_r \dots \#_r \xi''_k$$

with $\theta(\xi''_i) \equiv \langle b_i \rangle \equiv x_i \pmod{(\mu K)_r}$ for each i . Since $\theta(\xi') = x = x_1 \#_r \dots \#_r x_k$, it follows from Proposition 5.8 that $\theta(\xi''_i) = x_i$ for each i . But the x_i have unique inverse images ξ_i , so $\xi''_i = \xi_i$ for each i , and it follows that $\xi' = \xi$. This completes the proof. \square

The main theorem is now as follows.

Theorem 5.11. *The adjoint functors $\lambda: \omega\text{-cat} \rightarrow \mathbf{ADC}$ and $\nu: \mathbf{ADC} \rightarrow \omega\text{-cat}$ restrict to adjoint equivalences between the full subcategories consisting of ω -categories with loop-free atomic bases and of augmented directed complexes with loop-free unital bases. Under these equivalences, ω -categories with strongly loop-free atomic bases correspond to augmented directed complexes with strongly loop-free unital bases.*

Proof. By Theorem 5.6, if K is an augmented directed complex with a loop-free unital basis then νK has a loop-free atomic basis and the counit $\pi: \lambda \nu K \rightarrow K$ is an isomorphism.

Conversely, let C be an ω -category with a loop-free atomic basis E . By Definitions 4.4 and 4.5, the function $e \mapsto [e]_{|e|}$ maps E bijectively to a loop-free basis for λC . By Proposition 4.6 this basis is unital, so λC has a loop-free unital basis. By Proposition 4.6 again, the unit $\eta: C \rightarrow \nu \lambda C$ sends the generator e to the atom $\langle [e]_{|e|} \rangle$. It follows that η maps the generators of C bijectively to the atoms of $\nu \lambda C$ and preserves dimensions. By Theorem 5.7, $\eta: C \rightarrow \nu \lambda C$ is an isomorphism.

Finally, strongly loop-free atomic bases correspond to strongly loop-free unital bases by Definition 4.5(iii).

This completes the proof. \square

6. Relations with earlier work

In this section we compare our construction ν with other constructions of ω -categories.

We first observe that Theorem 5.11 serves to characterise certain ω -categories: if K is an augmented directed complex with a loop-free basis and if C is an ω -category with an atomic basis such that $\lambda C \cong K$, then C must be isomorphic to νK . This shows that the various constructions are essentially equivalent. For example, let C be the ω -category associated to the p -simplex in [2], [10] or [11]; then C must be isomorphic to $\nu\Delta[p]$, where $\Delta[p]$ is as in Example 3.8.

The ω -categories associated to loop-free structures in earlier treatments have presentations of particular types. The analogous result for the functor ν is as follows.

Theorem 6.1. *Let K be an augmented directed complex with a loop-free unital basis. Then the ω -category νK has a presentation as follows: the generators are the atoms; for each atom $\langle b \rangle$ there are relations $d_{|b|}^-(\langle b \rangle) = d_{|b|}^+(\langle b \rangle)$; for each positive-dimensional atom $\langle b \rangle$ there are relations $d_{|b|-1}^-(\langle b \rangle) = w^-(b)$ and $d_{|b|-1}^+(\langle b \rangle) = w^+(b)$, where the $w^\alpha(b)$ are expressions for the $d_{|b|-1}^\alpha(\langle b \rangle)$ as composites of atoms of dimension less than $|b|$.*

Proof. It follows from Theorem 5.5 that νK is composition-generated by its atoms, and the atoms therefore satisfy relations of the form described. Let C be the ω -category generated by the atoms subject to these relations; then there is a canonical morphism $\theta: C \rightarrow \nu K$, and we must show that it is an isomorphism. Because of Theorem 5.7, it suffices to show that the atoms composition-generate C . Because of Definition 2.1(iii) and (v), C is composition-generated by the elements $d_n^\alpha(\langle b \rangle)$ for $\langle b \rangle$ an atom. It therefore suffices to show that in C each element $d_n^\alpha(\langle b \rangle)$ is a composite of atoms of dimension at most n . We do this by induction on $|b|$. Indeed, if $n < |b|$ then $d_n^\alpha(\langle b \rangle) = d_n^\alpha d_{|b|-1}^\alpha(\langle b \rangle) = d_n^\alpha w^\alpha(b)$, and $w^\alpha(b)$ is a composite of atoms of dimension less than $|b|$, so the result holds by the inductive hypothesis and Definition 2.1(v), and if $n \geq |b|$ then $d_n^\alpha(\langle b \rangle) = d_n^\alpha d_{|b|}^\alpha(\langle b \rangle) = d_{|b|}^\alpha(\langle b \rangle) = \langle b \rangle$, so the result holds trivially. This completes the proof. \square

The constructions corresponding to ν in earlier treatments are described combinatorially in terms of sets rather than algebraically. We will now show that νK can be described combinatorially when K has a loop-free unital basis B , in the sense that the elements of νK are determined by subsets of B .

Theorem 6.2. *Let K be an augmented directed complex with a loop-free unital basis and let x be a member of νK .*

- (i) *If x is a sum of atoms then x is an atom.*
- (ii) *If $x \equiv m_1 \langle b_1 \rangle + \dots + m_k \langle b_k \rangle \pmod{(\mu K)_r}$ for some $r \geq -1$, with $|b_i| > r$ and with $m_i > 0$ for all i , then $m_i = 1$ for all i .*
- (iii) *Each term x_n^α of x is a sum of distinct basis elements.*

Proof. (i) Suppose that $x = \langle b_1 \rangle + \dots + \langle b_k \rangle$. Then $\varepsilon x_0^\alpha = \varepsilon \langle b_1 \rangle_0^\alpha + \dots + \varepsilon \langle b_k \rangle_0^\alpha$. But $\varepsilon x_0^\alpha = 1$ because $x \in \nu K$, and $\varepsilon \langle b_i \rangle_0^\alpha = 1$ for each i because the basis is unital, so $k = 1$. This means that x is equal to the atom $\langle b_1 \rangle$.

(ii) The proof is by induction on r . In the case $r = -1$ the result follows from part (i). From now on, suppose that $r \geq 0$. We may assume that $k \geq 1$ (otherwise there is nothing to prove). We have $\langle b_i \rangle_r^+ \wedge \langle b_i \rangle_r^- = 0$ for each i , because $|b_i| > r$. Since the basis is loop-free, we can assume the list $\langle b_1 \rangle, \dots, \langle b_k \rangle$ ordered so that $\langle b_i \rangle_r^+ \wedge \langle b_j \rangle_r^- = 0$ for $i > j$. By applying Proposition 5.1 and grouping the factors, we get a decomposition $x = x_1 \#_r \dots \#_r x_k$ with $x_i \in \nu K$ and $x_i \equiv m_i \langle b_i \rangle$ modulo $(\mu K)_r$. By Proposition 5.2(ii),

$$x_i \equiv m_i \langle b_i \rangle + \langle c_1 \rangle + \dots + \langle c_l \rangle \pmod{(\mu K)_{r-1}}$$

for some r -dimensional atoms $\langle c_1 \rangle, \dots, \langle c_l \rangle$. By the inductive hypothesis, $m_i = 1$ as required.

(iii) We have $x_n^\alpha \geq 0$ because $x \in \nu K$, so $x_n^\alpha = m_1 b_1 + \dots + m_k b_k$ for some positive integers m_i and some n -dimensional basis elements b_i . It now suffices to show that $m_i = 1$ for each i , and this follows from part (ii) because

$$d_n^\alpha x \equiv m_1 \langle b_1 \rangle + \dots + m_k \langle b_k \rangle \pmod{(\mu K)_{n-1}}.$$

□

7. The structure of the category of ω -categories

The ω -categories $F[p]$, $F[p; n]$, $F[p; n, n]$ and $F[p; n, m, n]$ of Example 4.7 represent the elements, operations and defining identities of ω -categories; in particular $F[p; n, n]$ and $F[p; n, m, n]$ are what one needs for the identities of Definition 2.1(vi) and (vii). One can therefore use these ω -categories to give a ‘globular’ description of ω -cat. There are also simplicial and cubical descriptions, using the ω -categories $\nu\Delta[p]$ and $\nu Q[p]$ coming from Examples 3.8 and 3.10; see [1], [2], [11]. All of these descriptions are based on ω -categories with strongly loop-free atomic bases. Because of Theorem 5.11, one can express these descriptions in terms of augmented directed complexes; thus the theory of ω -categories can be expressed in terms of chain complexes. We will now use the globular description to get results on monoidal structures. The main novelty is the functoriality; see Theorem 7.6 in particular.

We first make the following observation.

Theorem 7.1. *Every ω -category is the colimit of a small diagram of ω -categories with strongly loop-free atomic bases.*

Proof. Let C be an ω -category. By Proposition 2.3, C is the union of the increasing sequence of sub- ω -categories C_p , where

$$C_p = d_p^- C = d_p^+ C = \{ x \in C : d_p^- x = x \} = \{ x \in C : d_p^+ x = x \}.$$

The structure of C is given by the unary operations d_n^α and the not everywhere defined binary composition operations $\#_n$. Now the elements and $\#_n$ -composable pairs in C_p are represented by the ω -categories $F[p]$ and $F[p; n]$ of Example 4.7, and it follows that C is the colimit of a small diagram in which the objects are copies of the $F[p]$ and $F[p; n]$. Indeed, one needs a copy of $F[p]$ for each element of C_p ; one needs a copy of $F[p; n]$ for each $\#_n$ -composable pair in C_p ; and one

needs morphisms to determine the inclusions $C_{p-1} \rightarrow C_p$, the operations d_n^α , and the composition operations $\#_n$. The result now holds since, by Example 4.7, the $F[p]$ and $F[p; n]$ have strongly loop-free atomic bases. \square

From Example 3.10, the category of augmented directed complexes with strongly loop-free unital bases has a monoidal structure based on the usual tensor product of chain complexes. From Theorem 5.11 we immediately get the following consequence.

Proposition 7.2. *There is a monoidal structure on the full subcategory of $\omega\text{-cat}$ consisting of ω -categories with strongly loop-free atomic bases. It is equivalent via λ and ν to the monoidal structure on the full subcategory of **ADC** consisting of atomic directed complexes with strongly loop-free unital bases.*

We now use Theorem 7.1 to get a monoidal structure on the category of ω -categories.

Theorem 7.3. *There is a colimit-preserving functor*

$$(C, C') \mapsto C \otimes C' : \omega\text{-cat} \times \omega\text{-cat} \rightarrow \omega\text{-cat}$$

extending the tensor product on ω -categories with strongly loop-free atomic bases. This functor is unique up to natural equivalence, and it determines a monoidal structure on $\omega\text{-cat}$.

Proof. Because of Theorem 7.1, it makes sense to define the functor by

$$C \otimes C' = \text{colim}(F \otimes F'),$$

where the colimit is taken over the morphisms $F \rightarrow C$ and $F' \rightarrow C'$ whose domains have strongly loop-free atomic bases. It is clear that this functor preserves colimits and that it extends the tensor product on ω -categories with strongly loop-free atomic bases. It is also clear that these properties determine the functor up to natural equivalence. We get a monoidal structure on $\omega\text{-cat}$ because we are starting from a monoidal structure on the subcategory. \square

We now wish to show that the monoidal structure on $\omega\text{-cat}$ is biclosed. We will use the following result.

Theorem 7.4. *Let \mathbf{F} be the category of ω -categories with strongly loop-free atomic bases. Then an ω -category is naturally equivalent under the functor $C \mapsto \text{hom}(-, C)$ to a contravariant set-valued functor on \mathbf{F} which takes colimits in $\omega\text{-cat}$ to limits in the category of sets.*

Proof. If C is an ω -category, then $\text{hom}(-, C)$ is a contravariant set-valued functor on $\omega\text{-cat}$ taking colimits to limits. It follows that the restriction $\text{hom}(-, C)|_{\mathbf{F}}$ is a contravariant set-valued functor on \mathbf{F} taking colimits in $\omega\text{-cat}$ to limits in the category of sets. The axioms of ω -category theory (Definition 2.1) reduce to these properties of $\text{hom}(-, C)|_{\mathbf{F}}$: indeed it suffices to take the ω -categories $F[p]$, $F[p; n]$, $F[p; n, n]$ and $F[p; n, m, n]$ of Example 4.7, all of which are in \mathbf{F} . The result follows. \square

The category \mathbf{F} of Theorem 7.4 is equivalent under λ to a subcategory of \mathbf{ADC} by Theorem 5.11, so Theorem 7.4 yields a description of ω -categories in terms of chain complexes. Note in particular that λ takes colimits in $\omega\text{-cat}$ to colimits in \mathbf{ADC} because it is a left adjoint.

From Theorem 7.4, in the standard way, we get the following result.

Theorem 7.5. *The monoidal structure on $\omega\text{-cat}$ is biclosed.*

Proof. Given ω -categories C and D , we must find ω -categories $\text{HOM}(C, D)$ and $\text{HOM}'(C, D)$ such that there are natural equivalences

$$\text{hom}(-, \text{HOM}(C, D)) \cong \text{hom}(- \otimes C, D), \quad \text{hom}(-, \text{HOM}'(C, D)) \cong \text{hom}(C \otimes -, D).$$

But it follows from Theorem 7.3 that $\text{hom}(- \otimes C, D)$ is a contravariant set-valued functor on $\omega\text{-cat}$ taking colimits to limits, and it then follows from Theorem 7.4 that $\text{hom}(- \otimes C, D)$ is naturally equivalent to $\text{hom}(-, \text{HOM}(C, D))$ for some ω -category $\text{HOM}(C, D)$. The construction of $\text{HOM}'(C, D)$ is similar. \square

The category of chain complexes has a well-known closed symmetric monoidal structure, and one can check that this induces a biclosed monoidal structure on augmented directed complexes. Indeed, let K and L be augmented directed complexes. If $n > 0$ then

$$\text{HOM}(K, L)_n = \prod_m \text{hom}(K_m, L_{m+n}),$$

while $\text{HOM}(K, L)_0$ consists of the pairs $(f, \varepsilon f)$ such that $f \in \prod_m \text{hom}(K_m, L_m)$ is a chain map from K to L and εf is an integer with $(\varepsilon f)(\varepsilon x) = \varepsilon(fx)$ for $x \in K_0$. If $f \in \prod_m \text{hom}(K_m, L_{m+n})$, then we write f_m for the component of f in $\text{hom}(K_m, L_{m+n})$. The boundary on $\text{HOM}(K, L)_n$ is given by

$$(\partial f)_m = \begin{cases} \partial \circ f_m - (-1)^n f_{m-1} \circ \partial & \text{if } m > 0, \\ \partial \circ f_m & \text{if } m = 0, \end{cases}$$

and by $\varepsilon(\partial f) = 0$ in the case $n = 1$. The augmentation on $\text{HOM}(K, L)_0$ is given by $(f, \varepsilon f) \mapsto \varepsilon f$. The submonoid $\text{HOM}(K, L)_n^*$ is given by the elements f of $\prod_m \text{hom}(K_m, L_{m+n})$ such that $f_m(K_m^*) \subset L_{m+n}^*$ for all m . The definition of $\text{HOM}'(K, L)$ is the same as for $\text{HOM}(K, L)$, except that the boundary formula changes to

$$(\partial f)_m = \begin{cases} (-1)^m (\partial \circ f_m - f_{m-1} \circ \partial) & \text{if } m > 0, \\ \partial \circ f_m & \text{if } m = 0. \end{cases}$$

The biclosed monoidal structures on augmented directed complexes and ω -categories are related as follows.

Theorem 7.6. *Let K and L be augmented directed complexes such that K has a strongly loop-free unital basis. Then there are natural isomorphisms*

$$\nu \text{HOM}(K, L) \cong \text{HOM}(\nu K, \nu L), \quad \nu \text{HOM}'(K, L) \cong \text{HOM}'(\nu K, \nu L).$$

Proof. We give the proof for the first of these isomorphisms. Because of Theorem 7.4, it suffices to show that there are natural bijections

$$\mathrm{hom}(F, \nu \mathrm{HOM}(K, L)) \cong \mathrm{hom}(F, \mathrm{HOM}(\nu K, \nu L))$$

for F an ω -category with a strongly loop-free atomic basis. But

$$\mathrm{hom}(F, \nu \mathrm{HOM}(K, L)) \cong \mathrm{hom}(\lambda F, \mathrm{HOM}(K, L)) \cong \mathrm{hom}(\lambda F \otimes K, L)$$

and

$$\mathrm{hom}(F, \mathrm{HOM}(\nu K, \nu L)) \cong \mathrm{hom}(F \otimes \nu K, \nu L) \cong \mathrm{hom}(\lambda(F \otimes \nu K), L)$$

because λ is left adjoint to ν , and $\lambda F \otimes K \cong \lambda(F \otimes \nu K)$ by Proposition 7.2. The result follows. \square

References

- [1] F. A. Al-Agl, R. Brown and R. Steiner, Multiple categories: the equivalence of a globular and a cubical approach, *Adv. Math.* 170 (2002), no. 1, 71–118.
- [2] F. A. Al-Agl and R. Steiner, Nerves of multiple categories, *Proc. London Math. Soc.* (3) 66 (1993), no. 1, 92–128.
- [3] R. Brown and P. J. Higgins, Cubical abelian groups with connections are equivalent to chain complexes, *Homology Homotopy Appl.* 5(2003), no. 1, 49–52.
- [4] S. E. Crans and R. Steiner, Presentations of omega-categories by directed complexes, *J. Austral. Math. Soc. Ser. A* 63(1997), no. 1, 47–77.
- [5] P. Gaucher, Combinatorics of branchings in higher dimensional automata, *Theory Appl. Categ.* 8(2001), no. 12, 324–376.
- [6] M. Johnson, The combinatorics of n -categorical pasting, *J. Pure Appl. Algebra* 62(1989), no. 3, 211–225.
- [7] M. M. Kapranov and V. A. Voevodsky, Combinatorial-geometric aspects of polycategory theory: pasting schemes and higher Bruhat orders (list of results), *Cahiers Topologie Géom. Différentielle Catég.* 32(1991), no. 1, 11–27.
- [8] A. Patchkoria, Chain complexes of cancellative semimodules, *Bull. Georgian Acad. Sci.* 162(2000), no. 2, 206–208.
- [9] A. J. Power, An n -categorical pasting theorem, in Category theory (Como, 1990), 326–358, *Lecture Notes in Math.* 1488 Springer, Berlin, 1991.
- [10] R. Steiner, The algebra of directed complexes, *Appl. Categ. Structures* 1(1993), no. 3, 247–284.

- [11] R. Street, The algebra of oriented simplexes, *J. Pure Appl. Algebra* 49(1987), no. 3, 283–335.
- [12] R. Street, Parity complexes, *Cahiers Topologie Géom. Différentielle Catég.* 32(1991), no. 4, 315–343; Corrigenda, *Cahiers Topologie Géom. Différentielle Catég.* 35(1994), no. 4, 359–361.

This article may be accessed via WWW at <http://www.rmi.acnet.ge/hha/>
or by anonymous ftp at
[ftp://ftp.rmi.acnet.ge/pub/hha/volumes/2004/n1a12/v6n1a12.\(dvi,ps,pdf\)](ftp://ftp.rmi.acnet.ge/pub/hha/volumes/2004/n1a12/v6n1a12.(dvi,ps,pdf))

Richard Steiner r.steiner@maths.gla.ac.uk

Department of Mathematics,
University of Glasgow,
University Gardens, Glasgow,
Scotland G12 8QW